

Rev05  
Update: Jan,07,2025

## DATASHEET


# IL-5, Human(CHO-expressed)

Cat. No.: Z02981

## Product Introduction

Species	Human
Protein Construction	IL-5 (Ile20-Ser134) Accession # P05113
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	ED <sub>50</sub> < 1.0 ng/ml, measured in a cell proliferation assay using TF-1 cells, corresponding to a specific activity of > 1.0 × 10 <sup>6</sup> units/mg.
Expression System	CHO
Apparent Molecular Weight	29~35 kDa, on SDS-PAGE under non-reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

## Examples



ED<sub>50</sub> < 1 ng/ml, measured in a cell proliferation assay using TF-1 cells, corresponding to a specific activity of > 1 × 10<sup>6</sup> units/mg.

## Background

**Target Background :** Interleukin-5 (IL-5), produced by mast cells, T cells and eosinophils, is responsible for the activities attributed to eosinophil differentiating factor, B cell growth factor II and T cell-replacing factor (TRF). It can increase production and mobilization of eosinophils and CD34+ progenitors from the bone marrow. IL-5 plays an important role in inducing cell-mediated immunity against parasitic infections and certain tumors. IL-5 also promotes differentiation of basophils and primes them for histamine and leukotriene release.

**Synonyms :** IL5; EDF; BCDFII; TRF

**For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.**

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China